

KCL-TH-98-62

hep-th/9812133

February 1, 2008

On Harmonic Superspace

P.S. Howe

Department of Mathematics

King's College, London

Abstract

A short survey of some aspects of harmonic superspace is given. In particular, the $d = 3, N = 8$ scalar supermultiplet and the $d = 6, N = (2, 0)$ tensor multiplet are described as analytic superfields in appropriately defined harmonic superspaces.

1 Introduction

Victor Ogievetsky made many significant contributions to theoretical physics during a long and distinguished career often conducted in difficult circumstances. Perhaps his most interesting work, at least from this author's point of view, was the development of a deeper understanding of the geometry of supersymmetry. This began in the seventies with his and Sokatchev's interpretation of the geometry of $N = 1$ supergravity as the generalisation of chirality to the curved superspace context. Some years later he and his group, now expanded to include Galperin, Ivanov and Kalitzin as well as Sokatchev (GIKOS), succeeded in finding the most useful generalisation of chirality to the case of flat space $N = 2$ supersymmetry, namely $N = 2$ harmonic superspace. This new approach to the geometry of supersymmetry has turned out to have wide applications to both geometry and physics. In this note I shall briefly review flat harmonic superspace, its relation to twistor theory, its generalisation to curved superspace and its potentially interesting application to some on-shell supermultiplets which arise naturally in string theory in the context of the Maldacena conjecture.

2 Harmonic superspace

In extended supersymmetry flat harmonic superspaces [4] are superspaces of the form $M_H = M \times \mathbb{F}$ where M is the corresponding Minkowski superspace and $\mathbb{F}$ is a coset space of the internal symmetry group which is chosen to be a compact, complex manifold, in fact a flag manifold. For example, in $D = 4$, the internal symmetry group is $U(N)$ (or sometimes just $SU(N)$), the relevant flags sequences $V_{k_1} \subset V_{k_2} \subset \dots \subset \mathbb{C}^N$ of subspaces with dimension k_i , and the flag manifold is the space whose points correspond to such flags. In particular, the flags $V_p \subset V_{N-q} \subset \mathbb{C}^N$, $p+q \leq N$, define (N, p, q) harmonic superspace [9] for which $\mathbb{F} = H \backslash U(N)$, the isotropy group being $U(p) \times U(N - (p+q)) \times U(q)$. This family generalises in a natural way the harmonic superspaces for $N = 2$ and $N = 3$ supersymmetry introduced by GIKOS, which are respectively $(2, 1, 1)$ [4] and $(3, 1, 1)$ [5] harmonic superspace in the above notation.¹

The harmonic superspace approach of GIKOS emphasises the group theoretic aspects of fields defined on such spaces rather than the holomorphic aspects. For this reason it has become standard practice to work on the space $\hat{M}_H = M \times G$ (where $G = U(N)$ or whatever internal symmetry group is appropriate). This is equivalent to working on M_H provided that the fields are restricted to being equivariant with respect to the subgroup H . Such a field is a map $F : \hat{M}_H \rightarrow V$, where V is a representation space for the group H , such that $F(z, hu) = M(h)F(z, u)$, for all $u \in G$, $h \in H$ and $z \in M$, where $M(h)$ denotes the action of H on V . An equivariant field of this type defines in a natural way a section of a vector bundle E over M_H with typical fibre V and the two types of object are in one-to-one correspondence.

In $D = 4$ (N, p, q) harmonic superspace u is written in index notation as u_I^i where the i index is acted on by $U(N)$ and the I index by H , and the inverse u^{-1} is denoted u_i^I . The index I splits under H as (r, R, r') where $r = 1, \dots, p$, $R = p+1, \dots, N-q$, $r' = N-q+1, \dots, N$. The following derivatives can be defined on equivariant fields F ,

$$D_{\alpha r} = u_r^i D_{\alpha i}; \quad D_{\alpha}^{r'} = u_i^{r'} \bar{D}_{\alpha}^i \quad (1)$$

These derivatives anticommute and so allow the introduction of generalised chiral fields, or Grassmann-analytic (G -analytic) fields, F , which satisfy

$$D_{\alpha r} F = D_{\alpha}^{r'} F = 0 \quad (2)$$

Moreover, for any fixed values of the u 's the derivatives $\{D_{\alpha r}, D_{\alpha}^{r'}\}$ define a CR structure on M ; that is, they are basis vectors for an involutive subbundle $\bar{K}$ of the complexified tangent bundle T_c and their complex conjugates are linearly independent of them at any point in M .² The space $\mathbb{F}$ can thus be viewed as the space of all CR structures of this type on M . In addition, the derivatives can be combined with a subset of the right-invariant vector fields D_I^J on $U(N)$ to define a CR structure on M_H . The right-invariant vector fields decompose under H into the following subsets:

¹(3,2,1) superspace was first discussed in [6]

²The rôle of CR structures in harmonic superspace was first emphasised by [23].

$$D_I^J = \{D_r^s, D_R^S, D_{r'}^{s'}\}; \quad \{D_r^S, D_r^{s'}, D_R^{s'}\}; \quad \{D_R^s, D_{r'}^s, D_{r'}^S\} \quad (3)$$

The first subset corresponds to the isotropy group, the second to the components of the usual $\bar{\partial}$ operator on $\mathbb{F}$ and the third to its conjugate ∂ . The CR structure on M_H is then determined by the derivatives $\{D_{\alpha r}, D_{\alpha'}^{r'}, D_r^S, D_r^{s'}, D_R^s\}$. Since the D_I^J are characterised by

$$D_I^J u_K^k = \delta_K^J u_I^k; \quad D_I^J u_k^K = -\delta_I^K u_k^J \quad (4)$$

it is straightforward to verify that this set of derivatives does indeed specify a CR structure. Finally, when $p = q$, there is a real structure on harmonic superspace defined by $u \mapsto \epsilon u$ where

$$\epsilon = \begin{pmatrix} 0 & 0 & 1_p \\ 0 & 1_{N-2p} & 0 \\ -1_p & 0 & 0 \end{pmatrix} \quad (5)$$

This can be combined with complex conjugation to act on equivariant fields by

$$F(z, u) \mapsto \widetilde{F(z, u)} = \overline{F(z, \epsilon u)} \quad (6)$$

For fields transforming under certain representations of the isotropy group it is possible to demand that they be real with respect to this conjugation.

3 Relation to twistor theory

The above discussion is reminiscent of Euclidean twistor theory for $\mathbb{R}^4$ where $\mathbb{CP}^1$ parametrises the space of (anti)-self-dual complex structures on $\mathbb{R}^4$ and where the antipodal map of $\mathbb{CP}^1$ defines the real structure (as it does in (2,1,1) harmonic superspace) (see, for example [22, 24]). Indeed, particularly in cases where the appropriate symmetry group is the superconformal group, flat-space harmonic superspace theory can be viewed as an example of the group-theoretic approach to twistor theory [3]. The full power of this method only becomes apparent in complexified spacetime (or superspace), in which case the generalised twistor method starts with the selection of two parabolic subgroups P_1, P_2 of a complex simple group G . Since $P_{12} \equiv P_1 \cap P_2$ is also parabolic there are three (complex) homogeneous spaces $\mathbb{M}_k = P_k \backslash G$, $k = 1, 2$ or 12 which fit together into a double fibration $\mathbb{M}_2 \xleftarrow{\pi_2} \mathbb{M}_{12} \xrightarrow{\pi_1} \mathbb{M}_1$ such that each projection injects the fibres of the other one and thereby sets up a correspondence between points of $m_2 \in \mathbb{M}_2$ and subsets $\pi_1 \circ \pi^{-1}(m_2) \subset \mathbb{M}_1$ and vice versa. The basic idea of twistor theory applied to physics is that one of the spaces, say $\mathbb{M}_1$, is spacetime or Minkowski superspace and that information about field theory on (super) Minkowski space can be encoded as holomorphic data on $\mathbb{M}_2$. $N = 1$ supersymmetry in four dimensions provides an exceptional case where $\mathbb{M}_1$ and $\mathbb{M}_2$ are taken to be left and right chiral superspaces respectively and $\mathbb{M}_{12}$ is super Minkowski space [20].

In four dimensions the complex superconformal group is $SL(4|N)$. The parabolic isotropy groups define flag supermanifolds where the flags are now made of sequences of super subspaces of N -extended twistor space $\mathbb{T}_N = \mathbb{C}^{4|N}$, and the superspaces of interest are constructed as certain open subsets of these supermanifolds [20], see also [12]. (The flag supermanifolds themselves have compact bodies). As an example consider $N = 2$. The group $SL(4|2)$ consists of supermatrices of the form

$$g = \left(\begin{array}{c|c} A & \Gamma \\ \hline \Delta & B \end{array} \right) \quad (7)$$

where A and B are 4×4 and 2×2 even matrices, Γ and Δ are 4×2 and 2×4 respectively odd matrices, and where $\text{sdet} g = 1$. $N = 2$ super Minkowski space is the space of flags of type (i.e. dimension) $((2|0), (2|2))$ specified by the isotropy group consisting of matrices of the form

$$\left(\begin{array}{cccc|cc} \times & \times & & & & \\ \times & \times & & & & \\ \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \\ \hline \times & \times & & & \times & \times \\ \times & \times & & & \times & \times \end{array} \right) \quad (8)$$

where the crosses denote entries which are not necessarily zero. The body of this superspace can be identified from the top left part of the diagram which represents ordinary complex Minkowski space. Complex $N = 2$ harmonic superspace $\mathbb{M}_H$ is the space of flags of type $((2|2), (2|1), (2|2)) \subset \mathbb{T}_2$ specified by the subgroup of matrices of the form

$$\left(\begin{array}{cccc|cc} \times & \times & & & & \\ \times & \times & & & & \\ \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \\ \hline \times & \times & & & \times & \\ \times & \times & & & \times & \times \end{array} \right) \quad (9)$$

Note that this differs from super Minkowski space only in the bottom right-hand corner. Thus, $N = 2$ harmonic superspace has the same odd dimensionality as $N = 2$ super Minkowski space and its body is locally $\mathbb{M} \times \mathbb{CP}^1$. These two superspaces fit into a double fibration with analytic superspace $\mathbb{M}_A$ which the space of flags of type $(2|1) \subset \mathbb{T}_2$. The isotropy group consists of matrices of the form

$$\left(\begin{array}{cccc|cc} \times & \times & & & \times & \\ \times & \times & & & \times & \\ \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \\ \hline \times & \times & & & \times & \\ \times & \times & \times & \times & \times & \times \end{array} \right) \quad (10)$$

The odd dimensionality of analytic superspace is half that of harmonic superspace but its body is the same. The double fibration is $\mathbb{M}_A \xleftarrow{\pi_2} \mathbb{M}_H \xrightarrow{\pi_1} \mathbb{M}$. Points of $\mathbb{M}$ correspond to twistor lines, copies of the fibres of π_1 , in $\mathbb{M}_A$, while points of $\mathbb{M}_A$ parametrise certain planes of dimension $(0|4) \subset \mathbb{M}$, and these are copies of the fibres of π_2 . Note that analytic superspace is essentially a complex space; it does not exist as a coset space of the real $N = 2$ superconformal group $SU(2, 2|2)$. Furthermore, it is not a subspace of complex harmonic superspace, but rather a quotient space defined by the projection π_2 . The antipodal map on $\mathbb{CP}^1$ defines an anti-holomorphic involution of $\mathbb{M}_A$ and the real twistor lines in $\mathbb{M}_A$ are parametrised by points in real Minkowski superspace. When super Minkowski space is taken to be real, i.e. when $\mathbb{M}$ is replaced by M , real harmonic superspace $M_H = \pi_1^{-1}(M)$ and π_2 embeds M_H as a submanifold of $\mathbb{M}_A$. In the real case, therefore, the double fibration can be replaced by the single fibration $M_H \rightarrow M$ while holomorphic fields on $\mathbb{M}_A$ are replaced by CR -analytic fields on M_H [12].

The above discussion generalises in a natural way to (N, p, q) harmonic superspace [9, 12]. An entirely different family of supermanifolds can be constructed by using isotropy groups which are completely filled in the bottom right corner, but which have different structures to Minkowski space in the top left corner. Such superspaces, which are usually referred to as supertwistor spaces and which have conventional twistor spaces as bodies, have also been used in the context of supersymmetric field theory (see, for example, [25, 20]).

4 Curved harmonic superspace

The notion of harmonic superspace generalises to curved superspace relatively straightforwardly [7, 8, 9]. Perhaps the simplest geometries that can be discussed in this context are $D = 4$ superconformal geometries ($N \leq 4$) [10]. A superconformal structure on a real $(4|4N)$ supermanifold is a choice of odd tangent bundle F (rank $(0|4N)$) such that the Frobenius tensor, defined by evaluating the commutator of two odd vector fields (sections of F) modulo F has the following components with respect to bases $\{E_{\alpha i}, \bar{E}_{\dot{\alpha}}^i\}$ of F and $\{E^a\}$ for B^* where $B = T/F$:

$$T_{\alpha i \beta j}{}^c = T_{\dot{\alpha} \dot{\beta}}{}^{ij}{}^c = 0; \quad T_{\alpha i \dot{\beta}}{}^{jc} = -i \delta_i{}^j (\sigma^c)_{\alpha \dot{\beta}} \quad (11)$$

The Frobenius tensor is just the dimension zero component of the usual torsion tensor, although it is not necessary to introduce a connection to define it. With the components satisfying (11) above it is invariant under the group $\mathbb{R}^+ \times SL(2, \mathbb{C}) \cdot U(N)$ acting on F . The above constraints can be shown to lead to the usual constraints of off-shell conformal supergravity if standard conventional choices are made for the even tangent bundle B and for the $SL(2, \mathbb{C}) \cdot U(N)$ connection [10]. For $N = 4$ an additional constraint must be imposed arising from the fact that the true internal gauge symmetry group in this case is $SU(4)$.

In the curved case the space $\hat{M}_H$ is the (perhaps locally defined) principal $U(N)$ bundle associated with the $U(N)$ factor of the structure group of F , and harmonic superspace M_H itself is then given as $H \backslash \hat{M}_H$ where H is the relevant isotropy group, so that $\hat{M}_H$ is also a principal H -bundle over M_H .

For $N = 1$ harmonic superspace is not necessary and the constraints (11) above imply that a real $(4|4)$ supermanifold has a superconformal structure if and only if it has a CR structure with involutive CR bundle of rank $(0|2)$ such that the Frobenius tensor completely specifies B . This is Ogievetsky-Sokatchev supergeometry [21].

For $N = 2$, the appropriate harmonic superspace is $(2, 1, 1)$ superspace [8]. There is a natural bundle $\bar{K}$ on M_H defined by the set of basis vector fields D_1^2 and the horizontal lifts of the odd basis vector fields on M projected in the $(\alpha 1)$ and $\begin{pmatrix} 2 \\ \dot{\alpha} \end{pmatrix}$ directions. To define the latter it is necessary to introduce a $U(2)$ connection. This bundle defines a CR structure if and only if the constraints of $N = 2$ superconformal geometry are satisfied [9]. Note, however, that there is no gauge-invariant notion of G -analyticity. This problem was circumvented in [8] by allowing u_1^i and u_2^i to be independent, but the geometrical status of this strategy could perhaps be made more precise.

For $N = 3$ in $(3, 1, 1)$ harmonic superspace a similar CR structure is compatible with the constraints of conformal supergravity but does not imply them³, although demanding that the lift of the Frobenius tensor to M_H should vanish along the $(\alpha 1)$ and $\begin{pmatrix} 3 \\ \dot{\alpha} \end{pmatrix}$ directions does. In addition, at the linearised level, the conformal supergravity potential in harmonic superspace is a dimension -2 G -analytic field V_1^3 . This field can be interpreted as a 3-form along the antiholomorphic fibre directions, but a clear geometrical understanding is lacking at present.

Finally, for $N = 4$, it is possible to use $(4, 2, 2)$ harmonic superspace in which case the natural CR structure is equivalent to the constraints given above in (11). However, as noted previously, an additional constraint has to be imposed and this should eventually lead to the construction of a prepotential which is known to be a G -analytic 4-form along the antiholomorphic fibre directions in the linearised theory [9]. The case of $N = 4$ conformal supergravity is particularly interesting in view of its relevance to the Maldacena conjecture.

5 Superconformal fields

In many supersymmetric theories for which there is no known off-shell formulation the fundamental field may be considered to be a field strength tensor, and in many cases this field turns out to have a simple description as a single-component CR -analytic field W on an appropriate harmonic superspace. The hope is that this point of view may be useful in constructing, or attempting to construct, non-perturbative correlation functions of gauge-invariant composite operators made from powers of W by exploiting CR -analyticity, particularly when the theory under consideration is superconformal. This approach to quantum supersymmetry has been studied extensively in the case of $N = 4$ Yang-Mills theory (see, for example, [16], [17] and [15]) and is currently highly topical in that it is directly relevant to the Maldacena conjecture which relates supergravity theories on anti-de Sitter backgrounds to conformal field theories on the boundary, i.e. (super) Minkowski space [19].

The multiplets to be discussed here are matter (i.e. non-gravitational) multiplets with the maximal number of supersymmetries, sixteen. The basic multiplets are the $D = 4, N = 4$ Yang-Mills field strength multiplet, the $D = 6, (2, 0)$ tensor multiplet and the $D = 3, N = 8$ scalar multiplet. They are relevant to IIB string theory on $AdS_5 \times S^5$, M-theory on $AdS_7 \times S^4$ and M-theory on $AdS_4 \times S^7$ respectively, although the latter two cases are less well understood as these multiplets are only known at the non-interacting level. Nevertheless they are extremely interesting multiplets, corresponding as they do to the multiplets of the M-theory 5-brane and 2-brane respectively.

³This point has been observed by others (E. Sokatchev, private communication).

5.1 $D = 4, N = 4$ Yang-Mills

The field strength superfield $W_{ij} = -W_{ji}$ in super Minkowski space transforms according to the six-dimensional representation of the internal symmetry group $SU(4)$ and satisfies the following conditions:

$$\nabla_{\alpha i} W_{jk} = \nabla_{\alpha [i} W_{jk]} \quad (12)$$

$$\bar{\nabla}_{\dot{\alpha}}^i W_{jk} = -\frac{2}{3} \delta_{[j}^i \nabla_{\dot{\alpha}}^l W_{k]l} \quad (13)$$

$$W^{*ij} = \frac{1}{2} \epsilon^{ijkl} W_{kl} \quad (14)$$

The appropriate superspace in this case is $(4, 2, 2)$ harmonic superspace specified by the internal flag manifold $\mathbb{F} = S(U(2) \times U(2)) \backslash SU(4)$. An element $u \in SU(4)$ is written $u_I^i = (u_r^i, u_{r'}^i)$, $r = 1, 2$; $r' = 3, 4$. The right-invariant derivatives on $SU(4)$ are $\{D_r^s, D_{r'}^{s'}, D_o; D_r^{s'}, D_{r'}^s\}$, where $\{D_r^s, D_{r'}^{s'}, D_o\}$ correspond to the isotropy algebra $\mathfrak{su}(2) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)$, $D_r^{s'}$ corresponds to the $\bar{\partial}$ operator on $\mathbb{F}$ and $D_{r'}^s$ to its conjugate ∂ . The normalisation is

$$D_o u_r^i = \frac{1}{2} u_r^i; \quad D_o u_{r'}^i = -\frac{1}{2} u_{r'}^i \quad (15)$$

The CR -structure on harmonic superspace is specified by the set of derivatives $\{D_{\alpha r}, D_{\dot{\alpha}}^{r'}, D_r^{s'}\}$.

The claim is that W_{ij} is equivalent to a charge 1 field W on M_H which is covariantly G -analytic and ordinarily $\mathbb{F}$ -analytic; it is also real with respect to the real structure discussed in section 2 above, where covariantly G -analytic means that

$$\nabla_{\alpha r} W = \nabla_{\dot{\alpha}}^{r'} W = 0 \quad (16)$$

with $\nabla_{\alpha r} = u_r^i \nabla_{\alpha i}$, etc. This claim is easily verified [9].

The gauge-invariant operators $A_q = \text{tr}(W^q)$ are G -analytic in the usual sense and hence CR -analytic. These are the conformal fields of interest in the Maldacena conjecture. Since they are in short representations of $SU(2, 2|4)$, the integer q cannot be affected by quantum corrections and so, since this integer determines the dimension, they do not have anomalous dimensions. This family of operators was introduced in [16] and it has been shown that it is in one-to-one correspondence with the Kaluza-Klein spectrum of IIB supergravity on $AdS_5 \times S^5$ [1]. In particular, the family of operators includes the energy-momentum tensor $T = A_2$ [14]. In complex spacetime A_q becomes a holomorphic section of the q th power of a certain homogeneous line bundle $\mathcal{L}$ over analytic superspace. This superspace is in fact the super-Grassmannian whose points are $(2|2)$ planes in $\mathbb{T}_4 = \mathbb{C}^{4|4}$.

5.2 The $D = 6, (2, 0)$ tensor multiplet

In six-dimensional $(n, 0)$ supersymmetry the internal symmetry group is $Sp(n)$ and the spinors are taken to be symplectic Majorana-Weyl. The $(2, 0)$ tensor multiplet is a scalar superfield W_{ij} which transforms according to the real 5-dimensional representation of $Sp(2)$ (vector representation of $SO(5)$). It satisfies the constraints [13]

$$W_{ij} = -W_{ji}; \quad \eta^{ij} W_{ij} = 0 \quad (17)$$

$$\bar{W}^{ij} = \eta^{ik} \eta^{jl} W_{kl} \quad (18)$$

$$D_{\alpha i} W_{jk} = 2\eta_{ij} \lambda_{\alpha k} - 2\eta_{ik} \lambda_{\alpha j} + \eta_{jk} \lambda_{\alpha i} \quad (19)$$

where $\lambda_{\alpha i} = \frac{1}{5} \eta^{jk} D_{\alpha j} W_{ki}$, and where η_{ij} is the antisymmetric $Sp(2)$ invariant tensor. The components of W_{ij} consist of five scalars, four spinors and a self-dual three-form field strength tensor.

The appropriate harmonic superspace in this case is $M_H = M \times \mathbb{F}$ where $\mathbb{F} = U(2) \backslash Sp(2)$ [11]. Elements u of the group $Sp(2)$ are written $u_I^i = (u_r^i, u_{r'}^i)$, $r = 1, 2$; $r' = 3, 4$. In addition to being unitary the matrices u also satisfy $u_I^i u_J^j \eta_{ij} = \eta_{IJ}$. The matrix η has components

$$\eta_{rs} = \eta_{rs'} = 0; \quad \eta_{rs'} = -\eta_{s'r} = \delta_{rs'} \quad (20)$$

The right-invariant vector fields $\hat{D}_{IJ}$ on the group are real, symmetric and satisfy

$$\hat{D}_{IJ}u_K{}^i = -\eta_{IK}u_J{}^i - \eta_{JK}u_I{}^i \quad (21)$$

The derivatives split into the isotropy group derivatives $\hat{D}_{rs'}$ and the coset derivatives $\hat{D}_{rs} \equiv D_{rs}$ and $\hat{D}_{r's'} \equiv D_{r's'}$. The isotropy group derivatives may be separated into a $\mathfrak{u}(1)$ derivative $D_o = \frac{1}{2}\eta^{rs'}D_{rs'}$ and $\mathfrak{su}(2)$ derivatives $D_{rs'} = \hat{D}_{rs'} - \eta_{rs'}D_o$. On u , $D_o u_r{}^i = u_r{}^i$; $D_o u_{r'}{}^i = -u_{r'}{}^i$. The element $u_r{}^i$ transforms as a doublet under $\mathfrak{su}(2)$ while $u_{r'}{}^i$ transforms as the conjugate doublet.

The CR structure on M_H is specified by the derivatives $D_{\alpha r} = u_r{}^i D_{\alpha i}$ and D_{rs} , the latter corresponding to the components of $\bar{\partial}$ on $\mathbb{F}$ as usual.

In a very similar manner to the $D = 4, N = 4$ Yang-Mills case discussed above it can easily be shown that the field W_{ij} is equivalent to a single-component CR -analytic field W on M_H with $W = \frac{1}{2}\epsilon^{rs}u_r{}^i u_s{}^j W_{ij}$. It is also real with respect to a suitably defined real structure. Although there is no known non-Abelian version of this multiplet it is still possible to use it to discuss conformal field theory in the abstract [2]. The powers of W give conformal fields in short representations, in particular $W^2 = T$ is again the energy-momentum tensor multiplet (conformal supercurrent) [13]; a few of them were listed in [18]. It is likely that the complete set of such fields corresponds to the Kaluza-Klein spectrum of eleven-dimensional supergravity on $AdS_7 \times S^4$.

5.3 D=3, N=8 scalar multiplet

The scalar multiplet in three dimensions is slightly trickier to describe than the preceding two cases because the internal symmetry group is $SO(8)$ and it is necessary to use both spinor and vector representations of the group to describe the multiplet. Nevertheless it can be done and the result is simply stated: the $D = 3, N = 8$ scalar multiplet can be represented by a single-component CR -analytic superfield defined on the harmonic superspace $M_H = M \times U(4) \backslash Spin(8)$ [11].

The multiplet is defined in $D = 3, N = 8$ Minkowski space by the equation

$$D_A W_I = (\Sigma_I)_{AA'} \lambda^{A'} \quad (22)$$

where I, A and A' transform according to the vector, spinor and primed spinor representations of $Spin(8)$, Σ denotes the spin matrices and where the $D = 3$ spinor indices on both D and λ have been suppressed. The spinor representations decompose into 4 and $\bar{4}$ representations under $SU(4)$ with particular charges under $U(1)$ while the vector representation decomposes into the six-dimensional representation of $SU(4)$, neutral under $U(1)$ and a charged singlet and its conjugate under $U(1)$. That is $v_I \rightarrow (v_{++}, v_{--}, v_i)$, $i = 1 \dots 6$, where $v_{++} = \frac{1}{2}(v_7 - iv_8)$, $v_{--} = \bar{v}_{++}$, while for a $D = 8$ Majorana spinor Ψ , with

$$\Psi = \begin{pmatrix} \psi \\ \chi \end{pmatrix} \quad (23)$$

the indices are split as follows,

$$\psi_A \rightarrow \begin{pmatrix} \psi_{\alpha+} \\ \bar{\psi}_{\dot{\alpha}} \end{pmatrix} \quad \chi^{A'} \rightarrow \begin{pmatrix} \chi_{\alpha-} \\ \bar{\chi}_{\dot{\alpha}} \end{pmatrix} \quad (24)$$

The lower (upper) α indices correspond to the $4(\bar{4})$ representations of $SU(4)$.

The right-invariant derivatives on the group split into the isotropy derivatives $\{D_{ij}, D_o\}$ and the coset derivatives $\{D_{++i}, D_{--i}\}$. The CR structure is specified by the derivatives $D_{\alpha+} = u_{\alpha+}{}^A D_A$ and D_{++i} . The field $W_{++} = u_{++}{}^I W_I$ is easily seen to be CR -analytic, as well as being real with respect to an appropriate real structure.

In both of the latter two cases the superconformal group in complex spacetime is $OSp(8|2)$ and the analytic superspaces are the same, although the line bundles for the field strengths are different. However, the underlying twistor spaces are $\mathbb{C}^{4|8}$ for $D = 3$ and $\mathbb{C}^{8|4}$ for $D = 6$, the two being related by the Grassmann parity flip

operation Π . This may be a reflection of “electromagnetic” duality in $D = 11$; the 2-brane couples to the 4-form field strength of $D = 11$ supergravity while the 5-brane couples to its 7-form dual. In case (a) on the other hand, the D3-brane is self-dual, as is $D = 4, N = 4$ twistor space under Π .

References

- [1] L Andrianopoli and S Ferrara S (1998): *K-K excitations on $AdS_5 \times S^5$ and $N = 4$ primary superfields*, Phys. Lett. **B430** 248-253.
- [2] O Aharony, M Berkooz and N Seiberg (1998): *Light-cone description of $(2,0)$ conformal field theories in six dimensions*, Adv. Theor. Math. Phys. **2** 119-153.
- [3] R Baston and M Eastwood (1989): *The Penrose Transform*, (Oxford University Press).
- [4] A Galperin A, E Ivanov, S Kalitzin, V Ogievetsky and E Sokatchev (1984): *Unconstrained $N = 2$ matter, Yang-Mills and supergravity theories in harmonic superspace*, Class. Quantum Grav. **1** 469.
- [5] A Galperin, E Ivanov, S Kalitzin, V Ogievetsky and E Sokatchev (1985): *Unconstrained off-shell $N = 3$ supersymmetric Yang-Mills theory*, Class. Quantum Grav. **2** 155.
- [6] A Galperin A, E Ivanov, V Ogievetsky (1987): *Superspaces for $N = 3$ supersymmetry*, Soviet Journal for Nuclear Physics **46** 543.
- [7] A Galperin, Nguyen Ahn Ky and E Sokatchev (1987): *$N = 2$ supergravity: solution to the constraints*, Class. Quantum Grav. **4** 1235-1254.
- [8] A Galperin, E Ivanov, O Ogievetsky and E Sokatchev (1987): *$N = 2$ supergravity in superspace: different versions and matter couplings*, Class. Quantum Grav. **4** 1255.
- [9] G G Hartwell and P S Howe (1995): *(N, p, q) harmonic superspace*, Int J. Mod. Phys **10**, 3901-3919.
- [10] P S Howe (1981): *A superspace approach to conformal supergravity*, Phys. Lett. **100B** 389-392.
- [11] P S Howe (1998): *Superconformal fields in diverse dimensions*, in preparation.
- [12] P S Howe and G G Hartwell (1995): *A superspace survey*, Class. Quant. Grav. **12** 1823-1880.
- [13] P S Howe, G Sierra and P K Townsend (1983): *Supersymmetry in six dimensions* Nucl. Phys. **B221** (1983) 331-348.
- [14] P S Howe, K S Stelle and P K Townsend (1982): *Supercurrents*, Nucl. Phys. **B192** 332.
- [15] P S Howe P S, E Sokatchev and P C West (1998): *Three-point functions in $N = 4$ Yang-Mills*, hep-th/9808162.
- [16] P S Howe and P C West (1995): *Nonperturbative Green’s functions in theories with extended superconformal symmetry*, hep-th/9509140.
- [17] Howe P S, West P C (1995): *Superconformal invariants and extended supersymmetry*, Phys. Lett. **B400** 307-313.
- [18] R G Leigh and M Rozali (1998): *The large N limit of $(2,0)$ superconformal field theory*, Phys. Lett. **B431** 311-316.
- [19] J Maldacena J (1997): *The large N limit of superconformal field theories and supergravity*, hep-th/9711200.
- [20] Y I Manin (1988): *Gauge Field Theory and Complex Geometry* (Springer-Verlag).
- [21] V Ogievetsky and E Sokatchev (1980): *Soviet J. of Nucl. Phys.* **32** 447.
- [22] R Penrose R and W Rindler (1986): *Spinors and Spacetime II* (Cambridge University Press).
- [23] A A Rosly and A S Schwarz (1986): *Supersymmetry in spaces with auxiliary variables*, Comm. Math. Phys. **105**, (1986) 645.
- [24] R S Ward and R O Wells (1990): *Twistor Geometry and Field Theory* (Cambridge University Press).
- [25] E Witten E (1978): *Phys. Lett.* **77B**:394–398.